# CONTENTS

*JANUARY*


EDITORIALS

1 I: Can we characterize the central nervous system actions of $\alpha_2$-adrenergic agonists?
*M. E. P. Jones and M. Maze*

3 II: Publishing changes in 2001
*G. Smith*

CLINICAL INVESTIGATIONS

5 Sedative, analgesic, and cognitive effects of clonidine infusions in humans
*J. E. Hall, T. D. Uhrich and T. J. Ebert*

12 Model-based administration of inhalation anaesthesia. 1. Developing a system model
*J. G. C. Lerou and L. H. D. J. Booij*

29 Model-based administration of inhalation anaesthesia. 2. Exploring the system model
*J. G. C. Lerou and L. H. D. J. Booij*

38 Effects of sevoflurane and propofol on pulmonary shunt fraction during one-lung ventilation for thoracic surgery
*D. H. Beck, U. R. Doepfmer, C. Sinemus, A. Bloch, M. R. Schenk and W. J. Kox*

44 Recovery and neurological examination after remifentanil–desflurane or fentanyl–desflurane anaesthesia for carotid artery surgery
*W. Wilhelm, N. Schlaich, J. Harrer, S. Kleinschmidt, M. Müller and R. Larsen*

50 Comparison of changes in electroencephalographic measures during induction of general anaesthesia: influence of the gamma frequency band and electromyogram signal
*J. W. Sleigh, D. A. Steyn-Ross, M. L. Steyn-Ross, M. L. Williams and P. Smith*

59 Effects of different concentrations of sevoflurane and desflurane on subcortical somatosensory evoked responses in anaesthetized, non-stimulated patients
*D. J. A. Vaughan, C. Thornton, D. R. Wright, J. R. Fernandes, P. Robbins, C. Doré and M .D. Brunner*

63 Pre-existing cognitive impairment as a factor influencing outcome after cardiac surgery
*K. Millar, A. J. Asbury and G. D. Murray*

68 Fast-track cardiac anaesthesia in the elderly: effect of two different anaesthetic techniques on mental recovery
*N. P. Dowd, J. M. Karski, D. C. Cheng, S. Gajula, P. Seneviratne, J. A. Carroll Munro and D. Fiducia*

77 Intrathecal midazolam increases the analgesic effects of spinal blockade with bupivacaine in patients undergoing haemorrhoidectomy
*M. H. Kim and Y. M. Lee*

80 Comparison of two neurostimulation techniques for axillary brachial plexus blockade
*D. M. Coventry, K. F. Barker and M. Thomson*

84 Dexamethasone is a cost-effective alternative to ondansetron in preventing PONV after paediatric strabismus repair
*B. Subramaniam, R. Madan, S. Sadhasivam, B. Sennaraj, P. Tamilselvan, S. Rajeshwari, D. Jagan and D. Shende*

90 Comparison of effects of remifentanil and alfentanil on cardiovascular response to tracheal intubation in hypertensive patients
*A. M. Maguire, N. Kumar, J. L. Parker, D. J. Rowbotham and J. P. Thompson*

94 Comparison of coagulation and blood loss during anaesthesia with inhaled isoflurane or intravenous propofol
*N. L. Law, K. F. J. Ng, M. G. Irwin and J. S. F. Man*

99 Decontamination of laryngoscopes in The Netherlands
*M. J. L. Bucx, J. Dankert, M. M. Beenhakker and T. E. J. Harrison*

LABORATORY INVESTIGATION

103 Mechanical and electrophysiological effects of thiopental on rat cardiac left ventricular papillary muscle
*A. Descorps-Declère, M. P. Sauviat, K. Samii and Y. Lecarpentier*

REVIEW ARTICLE

110 Lazaroids: efficacy and mechanism of action of the 21-aminosteroids in neuroprotection
*R. J. Kavanagh and P. C. A. Kam*

SHORT COMMUNICATIONS

120 Pharmacokinetic profile of epidurally administered methylnaltrexone, a novel peripheral opioid antagonist in a rabbit model
*D. B. Murphy, H. El Behiery, V. W. Chan and J. F. Foss*

122 Influence of head and neck position on cuff position and oropharyngeal sealing pressure with the laryngeal mask airway in children
*K. Okuda, G. Inagawa, T. Miwa and K. Hiroki*

124 Waste gas exposure to sevoflurane and nitrous oxide during anaesthesia using the oesophageal–tracheal Combitube Small Adult™
*K. H. Hoerauf, T. Hartmann, S. Acimovic, A. Kopp, G. Wiesner, B. Gustorff, H. Jellinek and P. Krafft*

127 Effect of two anaesthetic regimens on airway nitric oxide production in horses
*D. J. Marlin, L. E. Young, R. McMurphy, K. Walsh and P. Dixon*

130 Use of cisatracurium during fast-track cardiac surgery
*A. Ouattara, L. Richard, J. M. Charrière, H. Lanquetot, P. Corbi and B. Debaene*

CASE REPORTS

133 Successful epidural anaesthesia for Caesarean section in a patient with spondyloepiphyseal dysplasia
*H. D. de Boer, A. Hemelaar, R. van Dongen and M. J. M. Gielen*

135 Use of remifentanil in a patient with peripartum cardiomyopathy requiring Caesarean section
*C. P. McCarroll, L. D. Paxton, P. Elliott and D. B. Wilson*

138 Reduction of vasopressor requirement by hydrocortisone administration in a patient with cerebral vasospasm
*J. A. Alhashemi*

141 Paradoxical changes in bispectral index during nitrous oxide administration
*G. D. Puri*

143 Therapeutic embolization of an internal mammary artery damaged by placement of a central venous catheter
*G. Abbott and P. Evans*

146 CORRESPONDENCE

157 BOOK REVIEWS

159 ERRATUM

*FEBRUARY*

EDITORIALS

161 I. Peri-operative β-blockade: a useful treatment that should be greeted with cautious enthusiasm
*S. J. Howell, J. W. Sear and P. Foex*

164 II. Is peri-operative renal dysfunction of no consequence?
*M. L. Cittanova*

166 III. A breakthrough in the genetic diagnosis of malignant hyperthermia
*R. L. Robinson and P. M. Hopkins*

CLINICAL INVESTIGATIONS

169 Haemodynamic and renal effects of intravenous enalaprilat during coronary artery bypass graft surgery in patients with ischaemic heart dysfunction
*F. Ryckwaert, P. Colson, J. Ribstein, G. Boccara and G. Guillon*

176 Is pulmonary resistance constant, within the range of tidal volume ventilation, in patients with ARDS?
*G. Mols, V. Kessler, A. Benzing, M. Lichtwarck-Aschoff, K. Geiger and J. Guttmann*

183 Gender differences in the pharmacokinetics of propofol in elderly patients during and after continuous infusion
*J. Vuyk, C. J. Oostwouder, A. A. Vletter, A. G. L. Burm and J. G. Bovill*

189 Comparing the effects of stimulation and propofol infusion rate on implicit and explicit memory formation
*J. Andrade, L. Englert, C. Harper and N. D. Edwards*

196 Absence of memory for intra-operative information during surgery with total intravenous anaesthesia
*I. F. Russell and M. Wang*

203 Psychomotor recovery in very old patients after total intravenous or balanced anaesthesia for cataract surgery
*J. Kubitz, J. Epple, A. Bach, J. Motsch, E. Martin and H. Schmidt*

209 EEG bispectral index and hypnotic component of anaesthesia induced by sevoflurane: comparison between children and adults
*C.-S. Degoute, C. Macabeo, C. Dubreuil, R. Duclaux and V. Banssillon*

213 Sevoflurane $EC_{50}$ and $EC_{95}$ values for laryngeal mask insertion and tracheal intubation in children
*R. Aantaa, R. Takala and P. Muittari*

217 Effect of common airway manoeuvres on upper airway dimensions and clinical signs in anaesthetized, spontaneously breathing children
*A. Reber, R. Paganoni and F. J. Frei*

223 Pharmacokinetics and pharmacodynamics of vecuronium in children receiving phenytoin or carbamazepine for chronic anticonvulsant therapy
*S. G. Soriano, L. J. Sullivan, K. Venkatakrishnan, D. J. Greenblatt and J. A. J. Martyn*

230 Epidural analgesia and arterial reconstructive surgery to the leg: effects on fibrinolysis and platelet degranulation
*S. A. Bew, A. E. Bryant, J. P. Desborough and G. M. Hall*

236 Intrathecal sufentanil and morphine for post-thoracotomy pain relief
*N. Mason, R. Gondret, A. Junca and F. Bonnet*

241 Spinal anaesthesia with ropivacaine 5 mg $ml^{-1}$ in glucose 10 mg $ml^{-1}$ or 50 mg $ml^{-1}$
*J. B. Whiteside, D. Burke and J. A. W. Wildsmith*

245 Rate of injection through Whitacre needles affects distribution of spinal anaesthesia
*L. Anderson, J. Walker, C. Brydon and M. G. Serpell*

LABORATORY INVESTIGATION

249 Comparison of relaxant effects of propofol on methacholine-induced bronchoconstriction in dogs with and without vagotomy
*S. Kabara, K. Hirota, E. Hashiba, H. Yoshioka, T. Kudo, T. Sato and A. Matsuki*

REVIEW ARTICLE

254 Anaesthesia and sleep apnoea
*J. A. Loadsman and D. R. Hillman*

SHORT COMMUNICATIONS

267 Renal effects of inhaled nitric oxide in humans
*W. M. Wraight and J. D. Young*

269 Can isoflurane mimic ischaemic preconditioning in isolated rat heart?
*N. Martini, B. Preckel, V. Thämer and W. Schlack*

272 Post-operative recovery: day surgery patients' preferences
*K. Jenkins, D. Grady, J. Wong, R. Correa, S. Armanious and F. Chung*

275 Attenuation of haemodynamic responses to tracheal intubation by the StyletScope
*T. Kitamura, Y. Yamada, M. Chinzei, H. L. Du and K. Hanaoka*

CASE REPORTS

278 Unexplained acute severe methaemoglobinaemia in a young adult
*M. Falkenhahn, S. Kannan and M. O'Kane*

280 Total spinal anaesthesia in association with insertion of a paravertebral catheter
*B. Lekhak, C. Bartley, I. D. Conacher and S. M. Nouraei*

COMMENTARY

283 Guidelines for molecular genetic detection of susceptibility to malignant hyperthermia
*A. Urwyler, T. Deufel, T. McCarthy and S. West for the European Malignant Hyperthermia Group*

288 CORRESPONDENCE

295 BOOK REVIEWS

298 ACKNOWLEDGEMENT OF ASSESSORS

301 ERRATA

ABSTRACTS

303P Proceedings of the Anaesthetic Research Society Meeting, London, November 16–17, 2000

*MARCH*

EDITORIALS

327 I. Right heart catheterization in intensive care
*J. D. Young*

329 II. Codeine phosphate in children: time for re-evaluation?
*M. Cunliffe*

CLINICAL INVESTIGATIONS

332 Dose-dependent prophylactic effect of nicorandil, an ATP-sensitive potassium channel opener, on intra-operative myocardial ischaemia in patients undergoing major abdominal surgery
*T. Kaneko, Y. Saito, Y. Hikawa, K. Yasuda and K. Makita*

338 Prediction of volatile anaesthetic solubility in blood and priming fluids for extracorporeal circulation
*R.-G. Yu, J.-X. Zhou and J. Liu*

345 Compound A production from sevoflurane is not less when KOH-free absorbent is used in a closed-circuit lung model system
*L. Versichelen, M. P. Bouche, M. Struys, J. Van Bocxlaer, E. Mortier, A. P. de Leenheer and G. Rolly*

349 Arterial to end-tidal carbon dioxide tension difference in children with congenital heart disease
*J. A. Short, S. T. Paris, P. D. Booker and R. Fletcher*

354 Biphasic EEG changes in relation to loss of consciousness during induction with thiopental, propofol, etomidate, midazolam or sevoflurane
*K. Kuizenga, J. M. K. H. Wierda and C. J. Kalkman*

361 Recovery from propofol anaesthesia supplemented with remifentanil
*R. A. O'Hare, R. K. Mirakhur, J. E. Reid, D. S. Breslin and A. Hayes*

366 Post-operative recovery after inguinal herniotomy in ex-premature infants: comparison between sevoflurane and spinal anaesthesia
*J. M. Williams, P. A. Stoddart, S. A. R. Williams and A. R. Wolf*

372 Evaluation of pre-emptive intramuscular phenylephrine and ephedrine for reduction of spinal anaesthesia-induced hypotension during Caesarean section
*B. T. Ayorinde, P. Buczkowski, J. Brown, J. Shah and D. J. Buggy*

377 I.V. ketoprofen for analgesia after tonsillectomy: comparison of pre- and post-operative administration
*A. Salonen, H. Kokki and K. Tuovinen*

382 Analysis of the French health ministry's national register of incidents involving medical devices in anaesthesia and intensive care
*L. Beydon, F. Conreux, R. Le Gall, D. Safran, J. B. Cazalaa and the members of the 'Sous-commission de Materiovigilance' for Anaesthesia and Intensive Care*

LABORATORY INVESTIGATIONS

388 Differential nitric oxide synthase activity, cofactor availability and cGMP accumulation in the central nervous system during anaesthesia
*H. F. Galley, A. E. Le Cras, S. D. Logan and N. R. Webster*

395 Suppressive effect of nitrous oxide on motor evoked potentials can be reversed by train stimulation in rabbits under ketamine/fentanyl anaesthesia, but not with additional propofol
*T. Sakamoto, M. Kawaguchi, S. Inoue and H. Furuya*

403 Evaluation and comparison of the pharmacokinetic and pharmacodynamic properties of allopregnanolone and pregnanolone at induction of anaesthesia in the male rat
*D. Zhu, M. D. Wang, T. Bäckström and G. Wahlström*

REVIEW ARTICLE

413 Codeine phosphate in paediatric medicine
*D. G. Williams, D. J. Hatch and R. F. Howard*

SHORT COMMUNICATIONS

422 Predicted effect compartment concentration of thiopental at loss of eyelash reflex
*T. A. Lim and K. Inbasegaran*

425 Effects of hypoxia and isoflurane on liver blood flow: role of adenosine
*S. H. Cray, M. W. Crawford, N. Khayyam and F. J. L. Carmichael*

427 Heterogeneous distribution of cardioplegic solution in pigs
*J.-S. Han, D.-M. Soh, C.-W. Joh, B.-I. Choi, Y.-S. Lee, C.-J. Lee and C.-H. Park*

CASE REPORTS

431 Liver transplantation for massive hepatic haemangiomatosis causing restrictive lung disease
*M. T. Keegan, G. S. Kamath, G. M. Vasdev, J. Y. Findlay, G. J. Gores, J. L. Steers and D. J. Plevak*

435 Undiagnosed cardiomyopathy in a neonate: significance of low oxygen saturation during anaesthesia
*W. Q. Smith and M. Abu-Harb*

437 Multiple acyl-CoA dehydrogenase deficiency: a rare cause of acidosis with an increased anion gap
*A. S. Grice and T. E. Peck*

442 Subarachnoid haemorrhage following spinal anaesthesia in an obstetric patient
*S. M. Eggert and K. A. Eggers*

COMMENTARY

445 Assessment instruments used during anaesthetic simulation: review of published studies
*A. J. Byrne and J. D. Greaves*

451 CORRESPONDENCE

460 BOOK REVIEWS

*APRIL*

EDITORIALS

465 I. KISS and indices of pulmonary oxygen transfer
*C. E. W. Hahn*

467 II. Continuous cardiac output by pulse contour analysis?
*J. J. van Lieshout and K. H. Wesseling*

469 III. Lessons from Stafford
*J. R. Sneyd*

473 IV. Local anaesthesia for eye surgery—no room for complacency
*G. S. Thind and A. P. Rubin*

CLINICAL INVESTIGATIONS

477 Effect of changes in arterial-mixed venous oxygen content difference ($C(a-\bar{v})_{O_2}$) on indices of pulmonary oxygen transfer in a model ARDS lung
*M. Nirmalan, T. Willard, M. O. Columb and P. Nightingale*

486 Estimation of changes in cardiac output from the arterial blood pressure waveform in the upper limb
*N. W. F. Linton and R. A. F. Linton*

497 Transoesophageal echocardiography in cardiac and vascular surgery: implications and observer variability
*D. Schmidlin, D. Bettex, E. Bernard, R. Germann, M. Tornic, R. Jenni and E. R. Schmid*

506 Intercurrent drug therapy and perioperative cardiovascular mortality in elective and urgent/emergency surgical patients
*J. W. Sear, S. J. Howell, Y. M. Sear, D. Yeates, M. Goldacre and P. Foex*

513 The auditory evoked response as an awareness monitor during anaesthesia
*E. Loveman, J. C. Van Hooff and D. C. Smith*

519 Ocular microtremor: a tool for measuring depth of anaesthesia?
*S. Bojanic, T. Simpson and C. Bolger*

523 Effects of fentanyl, alfentanil, remifentanil and sufentanil on loss of consciousness and bispectral index during propofol induction of anaesthesia
*C. Lysakowski, L. Dumont, M. Pellégrini, F. Clergue and E. Tassonyi*

528 Antinociceptive properties of neurosteroids IV: pilot study demonstrating the analgesic effects of alphadolone administered orally to humans
*C. S. Goodchild, A. Robinson and R. Nadeson*

535 Persistent back pain after spinal anaesthesia in the non-obstetric setting: incidence and predisposing factors
*K. Schwabe and H.-B. Hopf*

LABORATORY INVESTIGATIONS

540 Cytochrome P4502B6 and 2C9 do not metabolize midazolam: kinetic analysis and inhibition study with monoclonal antibodies
*N. Hamaoka, Y. Oda, I. Hase and A. Asada*

545 Naloxone improves functional recovery of myocardial stunning in conscious dogs through its action on the central nervous system
*T. P. Weber, J. Stypmann, A. Meißner, M. Große Hartlage, H. Van Aken and N. Rolf*

550 Effects of sevoflurane on dopamine, glutamate and aspartate release in an *in vitro* model of cerebral ischaemia
*C. C. Toner, K. Connelly, R. Whelpton, S. Bains, A. T. Michael-Titus, D. P. McLaughlin and J. A. Stamford*

REVIEW ARTICLE

555 Neurofibromatosis: clinical presentations and anaesthetic implications
*N. P. Hirsch, A. Murphy and J. J. Radcliffe*

SHORT COMMUNICATIONS

565 Intrapartum epidural catheter migration: a comparative study of three dressing applications
*S. M. Burns, C. M. Cowan, P. M. Barclay and R. G. Wilkes*

567 Choice of opioid for initiation of combined spinal epidural analgesia in labour—fentanyl or diamorphine
*D. J. A. Vaughan, N. Ahmad, N. K. Lillywhite, N. Lewis, D. Thomas and P. N. Robinson*

570 Small-dose selective spinal anaesthesia for short-duration outpatient gynaecological laparoscopy: recovery characteristics compared with propofol anaesthesia
*A. V. G. Stewart, H. Vaghadia, L. Collins and G. W. E. Mitchell*

572 Analgesic efficacy of tramadol 2 mg $kg^{-1}$ for paediatric day-case adenoidectomy
*H. Viitanen and P. Annila*

575 Reduced haemostatic factor transfusion using heparinase-modified thrombelastography during cardiopulmonary bypass
*D. Royston and S. von Kier*

578 Circulating tetrahydrobiopterin concentrations in patients with septic shock
*H. F. Galley, A. E. Le Cras, K. Yassen, I. S. Grant and N. R. Webster*

CASE REPORTS

581 Review of management of purpura fulminans and two case reports
*J. Nolan and R. Sinclair*

586 Intra-operative and post-operative hypercapnia leading to delayed respiratory failure associated with transanal endoscopic microsurgery under general anaesthesia
*K. Kerr and G. H. Mills*

589 Acute respiratory and metabolic acidosis induced by excessive muscle contraction during spinal evoked stimulation
*Y. Tohdoh, S. Sumita, T. Kawamata, K. Omote, S. Kawana and A. Namiki*

594 CORRESPONDENCE

603 BOOK REVIEWS

605 ERRATA


*MAY*


EDITORIALS

607 I. Mapping the anatomy of unconsciousness—imaging anaesthetic action in the brain
*D. K. Menon*

611 II. How can we improve the outcome of oesophagectomy?
*K. M. Sherry*

614 III. Development of the concept of a liquid pulmonary alveolar lining layer
*K. L. Dorrington and J. D. Young*

CLINICAL INVESTIGATIONS

618 Correlating *in vivo* anaesthetic effects with *ex vivo* receptor density data supports a GABAergic mechanism of action for propofol, but not for isoflurane
*M. T. Alkire and R. J. Haier*

627 Clonidine decreases propofol requirements during anaesthesia: effect on bispectral index
*S. B. Fehr, M. P. Zalunardo, B. Seifert, K. M. Rentsch, R. G. Rohling, T. Pasch and D. R. Spahn*

633 Peri-operative risk factors for acute lung injury after elective oesophagectomy
*S. Tandon, A. Batchelor, R. Bullock, A. Gascoigne, M. Griffin, N. Hayes, J. Hing, I. Shaw, I. Warnell and S. V. Baudouin*

639 Comparison of pattern of breathing with other measures of induction of anaesthesia, using propofol, methohexital, and sevoflurane
*T. L. Strickland and G. B. Drummond*

645 Uptake of isoflurane during prolonged clinical anaesthesia
*S. K. Pal, G. G. Lockwood and D. C. White*

650 Effects of dexmedetomidine on adrenocortical function, and the cardiovascular, endocrine and inflammatory responses in post-operative patients needing sedation in the intensive care unit
*R. M. Venn, A. Bryant, G. M. Hall and R. M. Grounds*

657 Does the optimization of cardiac output by fluid loading increase splanchnic blood flow?
*S. G. Sakka, K. Reinhart and A. Meier-Hellmann*

663 Amrinone can accelerate the cooling rate of core temperature during deliberate mild hypothermia for neurosurgical procedures
*S. Inoue, M. Kawaguchi, T. Sakamoto, T. Iwata, Y. Kawaraguchi, H. Furuya and T. Sakaki*

669 Autologous blood transfusion in total knee replacement surgery
*D. Thomas, K. Wareham, D. Cohen and H. Hutchings*

674 Double-blind comparison of ropivacaine 7.5 mg $ml^{-1}$ with bupivacaine 5 mg $ml^{-1}$ for sciatic nerve block
*C. Connolly, D. M. Coventry and J. A. W. Wildsmith*

678 Rocuronium: high risk for anaphylaxis?
*M. Rose and M. Fisher*

LABORATORY INVESTIGATIONS

683 Effects of haemoglobin-based oxygen carrier Hemoglobin glutamer-200 (bovine) on intestinal perfusion and oxygenation in a canine hypovolaemia model
*B. Driessen, J. S. Jahr, F. Lurie, S. M. Griffey and R. A. Gunther*

693 Tissue/blood and tissue/water partition coefficients for propofol in sheep
*B. M. Q. Weaver, G. E. Staddon and W. W. Mapleson*

704 Antinociceptive properties of neurosteroids III: experiments with alphadolone given intravenously, intraperitoneally, and intragastrically
*R. Nadeson and C. S. Goodchild*

REVIEW ARTICLE

709 Hormone replacement therapy (HRT) and anaesthesia
*D. Brighouse*

CASE REPORTS

717 Abnormal pupillary activity in a brainstem-dead patient
*D. Shlugman, M. Parulekar, J. S. Elston and A. Farmery*

720 Subdural haematoma after dural puncture headache treated by epidural blood patch
*J. M. Davies, A. Murphy, M. Smith and G. O'Sullivan*

723 Incremental spinal anaesthesia for elective Caesarean section in a patient with Eisenmenger's syndrome
*P. J. Cole, M. H. Cross and M. Dresner*

727 Undiagnosed myasthenia gravis unmasked by neuromuscular blockade
*M. F. Dunsire, S. G. Clarke and J. J. Stedmon*

731 Haemodynamic changes during retroperitoneoscopic adrenalectomy for phaeochromocytoma
*F. Atallah, T. Bastide-Heulin, M. Soulié, F. Crouzil, A. Galiana, K. Samii and C. Virenque*

734 CORRESPONDENCE

739 BOOK REVIEWS

742 ERRATUM

*JUNE*

EDITORIALS

743 I. Circulating volume and clinical assessment of the circulation
*M. Palazzo*

746 II. A momentary pause . . .
*M. A. Gropper*

749 III. Is it safe to artificially ventilate a paralysed patient through the laryngeal mask? The jury is still out
*G. Sidaras and J. M. Hunter*

CLINICAL INVESTIGATIONS

754 Clinical evaluation of circulating blood volume in critically ill patients—contribution of a clinical scoring system
*F. Stéphan, A. Flahault, N. Dieudonné, J. Hollande, F. Paillard and F. Bonnet*

763 Recovery after remifentanil and sufentanil for analgesia and sedation of mechanically ventilated patients after trauma or major surgery
*S. Soltész, A. Biedler, M. Silomon, I. Schöpflin and G. P. Molter*

769 Monitoring level of sedation with bispectral EEG analysis: comparison between hypothermic and normothermic cardiopulmonary bypass
*D. Schmidlin, P. Hager and E. R. Schmid*

777 Inspiratory timing during anaesthesia: a model of cardioventilatory coupling
*D. C. Galletly and P. D. Larsen*

789 Survey of cancellation rate of hypertensive patients undergoing anaesthesia and elective surgery
*P. Dix and S. Howell*

794 Oesophageal Doppler monitoring overestimates cardiac output during lumbar epidural anaesthesia
*H. A. Leather and P. F. Wouters*

798 The lumbar epidural space in pregnancy: visualization by ultrasonography
*T. Grau, R. W. Leipold, J. Horter, R. Conradi, E. Martin and J. Motsch*

805 Spinal anaesthesia for Caesarean section with bupivacaine 5 mg $ml^{-1}$ in glucose 8 or 80 mg $ml^{-1}$
*C. Connolly, G. A. McLeod and J. A. W. Wildsmith*

808 Does prior administration of enoxaparin influence the effects of levobupivacaine on blood clotting? Assessment using the Thrombelastograph®
*S. A. Leonard, A. Lydon, M. Walsh, C. Fleming, J. Boylan and G. D. Shorten*

814 Infusion of amino acid enriched solution hastens recovery from neuromuscular block caused by vecuronium
*Y. Saitoh, K. Kaneda, Y. Tokunaga and M. Murakawa*

822 What can be expected from risk scores for predicting postoperative nausea and vomiting?
*C. C. Apfel, P. Kranke, C.-A. Greim and N. Roewer*

828 Downward movement of syringe pumps reduces syringe output
*H. Kern, A. Kuring, U. Redlich, U. R. Döpfmer, N. M. Sims, C. D. Spies and W. J. Kox*

LABORATORY INVESTIGATIONS

832 Influence of sepsis on sevoflurane minimum alveolar concentration in a porcine model
*B. Allaouchiche, F. Duflo, J.-P. Tournadre, R. Debon and D. Chassard*

837 Halothane potentiates effect of methamphetamine and nomifensine on extracellular dopamine levels in rat striatum: a microdialysis study
*Y. U. Adachi, K. Watanabe, T. Satoh and E. S. Vizi*

846 Lidocaine reduces ischaemic but not reperfusion injury in isolated rat heart
*D. Ebel, P. Lipfert, J. Fräßdorf, B. Preckel, J. Müllenheim, V. Thämer and W. Schlack*

853 Propofol inhibits FMLP-stimulated phosphorylation of p42 mitogen-activated protein kinase and chemotaxis in human neutrophils
*T. Nagata, M. Kansha, K. Irita and S. Takahashi*

REVIEW ARTICLE

859 Perioperative bradycardia and asystole: relationship to vasovagal syncope and the Bezold–Jarisch reflex
*S. M. Kinsella and J. P. Tuckey*

SHORT COMMUNICATIONS

869 Gastric residual volume in children: a study comparing efficiency of erythromycin and metoclopramide as prokinetic agents
*T. F. Zatman, J. E. Hall and M. Harmer*

871 Effect of intravenous magnesium on pain and secondary hyperalgesia associated with the heat/capsaicin sensitization model in healthy volunteers
*S. Mikkelsen, J. Dirks, P. Fabricius, K. L. Petersen, M. C. Rowbotham and J. B. Dahl*

874 Orally administered clonidine significantly reduces pain during injection of propofol
*T. Yoshikawa, Z. Wajima, A. Ogura, T. Inoue and R. Ogawa*

876 Efficacy of varying concentrations of hyaluronidase in peribulbar anaesthesia
*C. Mantovani, A. E. Bryant and G. Nicholson*

879 Midazolam *versus* propofol for reducing contractility in fatigued canine diaphragm
*Y. Fujii and H. Toyooka*

CASE REPORTS

882 Xenon anaesthesia for laparoscopic cholecystectomy in a patient with Eisenmenger's syndrome
*J. Hofland, I. Gültuna and R. Tenbrinck*

886 Respiratory failure after tracheal extubation in a patient with halo frame cervical spine immobilization—rescue therapy using the Combitube airway
*M. Mercer*

891 Intramuscular ketamine in a parturient in whom pre-operative intravenous access was not possible
*W. C. Lum Hee and V. F. Metias*

893 Cranial subdural haematoma after spinal anaesthesia
*R. Acharya, S. S. Chhabra, M. Ratra and A. D. Sehgal*

896 CORRESPONDENCE

905 BOOK REVIEWS

909 INDEX TO VOLUME 86